

Epidemiologic Profile for HIV/AIDS in Indiana 2003

May 18, 2004



Acknowledgments

This profile was compiled by Marc Wiehn, Luther Consulting, LLC. Feedback and comments are welcome. Please forward comments to:

Marc V. Wiehn
Senior Associate – Evaluation Services
Luther Consulting, LLC
423 Massachusetts Avenue
Indianapolis, IN 46204
Email: mwiehn@evaluationweb.com
Phone: (317) 636-0282 Ext. 104

Indiana State Department of Health:

Jerry Burkman
Larry Harris
Shawn Carney
Linda Stemnock
Sextus Tansinin
Andrew Mitsch
Kevin Langford

Luther Consulting:

James Luther, PhD
John Roach
Zach Steiner
Reuben Craft
Anne Horowitz

Table of Contents

List of Figures and Tables

Executive Summary

Introduction

Introduction

Profile Data Sources

Strengths and Limitations of Profile

Organization of the Profile

Core Epidemiologic Questions

What are the sociodemographic characteristics of the general population?

What is the scope of the HIV/AIDS Epidemic in Indiana?

Ryan White CARE Act Questions

What are the patterns of service utilization of HIV-infected people?

Appendices

Appendix A: Data Sources

Appendix B: Glossary of Terminology and Acronyms

Appendix C: Map of Indiana with Health Regions

List of Figures and Tables

| Figures | | Page |
|-----------|-------------------------------------------------------------------------------|------|
| Figure 1 | Indiana's Urban and Rural Areas, 2000 | 20 |
| Figure 2 | Indiana's Most Populous Counties | 21 |
| Figure 3 | Percent Change in Indiana's Population, 2000-2010 | 22 |
| Figure 4 | Indiana Population by Race and Ethnicity, Census Estimates 2002 | 23 |
| Figure 5 | Indiana Population by Ethnicity, Census Estimates 2002 | 24 |
| Figure 6 | Racial and Ethnic Growth Rates for the past Decade | 24 |
| Figure 7 | Age Distribution of the Indiana Population, Census Bureau 2000 | 25 |
| Figure 8 | Population Changes by Age Groups | 26 |
| Figure 9 | Health Regions in Indiana | 27 |
| Figure 10 | Type of Housing in Indiana, Census 2000 | 28 |
| Figure 11 | Poverty Levels by Health Regions in Indiana, 2001 | 30 |
| Figure 12 | Prevalence Rates for HIV/AIDS by Sex, July 2002 to June 2003 | 36 |
| Figure 13 | Prevalence Rate for HIV/AIDS by Age, July 2002 to June 2003 | 37 |
| Figure 14 | Number of Persons infected with HIV and AIDS by Age at Diagnosis | 38 |
| Figure 15 | Prevalence Rates of HIV/AIDS Infected Persons by Race/Ethnicity in 2003 | 39 |
| Figure 16 | Prevalence Numbers in Percent of Total Infected by Race/Ethnicity and Sex | 40 |
| Figure 17 | Prevalence Rates for HIV/AIDS by Mode of Transmission, July 2002 to June 2003 | 42 |
| Figure 18 | Prevalence Rate for HIV/AIDS by Mode of Transmission and Race/Ethnicity | 43 |
| Figure 19 | Prevalence Numbers for HIV and AIDS by Heterosexual Contact Categories | 44 |
| Figure 20 | Prevalence Rate for HIV/AIDS by Health Region, July 2002 to June 2003 | 45 |
| Figure 21 | Prevalence for HIV and AIDS by Health Region, July 2002 to June 2003 | 46 |
| Figure 22 | Incidence Numbers of HIV and AIDS for Indiana, 1982 to 2002 | 49 |
| Figure 23 | New Infections (Incidence) Rate for HIV/AIDS by Sex, July 2002 to June 2003 | 52 |
| Figure 24 | Incidence Rate for HIV and AIDS by Sex, July 2002 to June 2003 | 52 |
| Figure 25 | Incidence Rate for HIV/AIDS by Age, July 2002 to June 2003 | 53 |

| | | |
|-----------|-----------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 26 | Incidence Rate for HIV and AIDS by Age, July 2002 to June 2003 | 54 |
| Figure 27 | Incidence Rate of HIV/AIDS by Race/Ethnicity, July 2002 to June 2003 | 55 |
| Figure 28 | Incidence Rate for HIV and AIDS by Race/Ethnicity, July 2002 to June 2003 | 56 |
| Figure 29 | Incidence Rates for HIV/AIDS by Race/Ethnicity by Sex, July 2002 to June 2003 | 57 |
| Figure 30 | Incidence Rates per 100,000 for HIV/AIDS by Mode of Transmission | 58 |
| Figure 31 | Incidence Rate of HIV/AIDS by Mode of Transmission and by Race/Ethnicity | 59 |
| Figure 32 | Incidence Rate/100,000 by Indiana Health Region, July 2002 to June 2003 | 61 |
| Figure 33 | Incidence Rates for HIV and AIDS by Health Region | 63 |
| Figure 34 | Time of Diagnosis of Seroconversion for Mothers in Indiana, June 2003 | 64 |
| Figure 35 | Number of HIV+ Mothers by Race/Ethnicity in June 2003 | 65 |
| Figure 36 | Cumulative Number of Children born to HIV Positive Mothers, June 2003 | 65 |
| Figure 37 | Infection Status of Children of HIV+ Mothers by the Time of HIV Detection, June 2003 | 66 |
| Figure 38 | Number of Deaths by Year in Indiana, 1981 to 2002 | 69 |
| Figure 39 | Mortality Rates by Age in Indiana, 2002 | 69 |
| Figure 40 | Mortality Rate by Sex for Indiana, 2002 | 70 |
| Figure 41 | Mortality Rate for Indiana by Race and Ethnicity, 2002 | 70 |
| Figure 42 | Mortality Rates by Mode of Transmission for Indiana, 2002 | 71 |
| Figure 43 | Mortality Rate by Health Regions for Indiana, 2002 | 72 |
| Figure 44 | Cumulative Number of Migrants by Sex | 75 |
| Figure 45 | Cumulative Percentage Rate of Migrants by Race and Ethnicity, 2003 | 76 |
| Figure 46 | Cumulative Number of Infected Persons Moving To Indiana by Health Region, 2003 | 77 |
| Figure 47 | Percentage of Migrants by Mode of Transmission, 2003 | 78 |
| Figure 48 | CTR Test Performed by Age Group in Indiana | 80 |
| Figure 49 | BRFSS Result: Have you ever been tested for HIV? | 84 |
| Figure 50 | BRFSS Result: Can a pregnant woman with HIV get treatment to help reduce the chances that she will pass the virus on to her baby? | 85 |
| Figure 51 | BRFSS Result: Are there medical treatments available that are intended to help a person who is infected with HIV to live longer? | 86 |
| Figure 52 | BRFSS Result: In the past 12 months has a health professional talked to you about preventing STD's through condom use? | 86 |

| | | |
|-----------|------------------------------------------------------------------------------------------------|-----|
| Figure 53 | STD Cases in Indiana, 1998-2003 | 87 |
| Figure 54 | STD Cases in Indiana by Sex, 2003 | 88 |
| Figure 55 | STD Cases in Indiana by Age, 2003 | 89 |
| Figure 56 | STD Cases in Indiana by Race, 2003 | 89 |
| Figure 57 | STD Cases in Indiana by Ethnicity, 2003 | 90 |
| Figure 58 | Number of Pregnancies in Indiana by Race, 2001 | 93 |
| Figure 59 | Number of Pregnancies in Indiana by Age and Race, 2001 | 93 |
| Figure 60 | Number of Terminated Pregnancies in Indiana by Age and Race, 2001 | 96 |
| Figure 61 | Percentage of Terminated Pregnancies in Indiana by Ethnicity, 2001 | 96 |
| Figure 62 | Number of ADAP Recipients by Year and Sex, Indiana | 102 |
| Figure 63 | Number of HIAP Recipients by Year and Sex, Indiana | 102 |
| Figure 64 | Relative Number of ADAP Recipients by Year and Race, 1999-2002 | 103 |
| Figure 65 | Relative Number of HIAP Recipients by Year and Race, 1999-2002 | 104 |
| Figure 66 | Relative Number of ADAP Recipients by Year and Ethnicity, 1999-2002 | 105 |
| Figure 67 | Relative Number of HIAP Recipients by Year and Ethnicity, 1999-2002 | 106 |
| Figure 68 | Number of ADAP and HIAP Recipients by Age, 2002 | 107 |
| Figure 69 | Number of Patients with No Reported Viral Load or CD4+ Count in 2002 by Race/Ethnicity | 109 |
| Figure 70 | Number of Patients with No Reported Viral Load or CD4+ Count in 2002 by Race/Ethnicity and Sex | 110 |

| Tables | | Page |
|---------------|----------------------------------------------------------------------------------------------------------|-------------|
| Table 1 | Estimated Population Size for Health Regions by Race/Ethnicity in 2002 | 28 |
| Table 2 | Poverty Level in Indiana by County of Residence, 2001 | 29 |
| Table 3 | Employment Type by County in Indiana, 2001 | 31 |
| Table 4 | Percentage distribution of adults (19-64) by Insurance Coverage and Gender, Indiana, 2000-2001 | 32 |
| Table 5 | Cumulative Number of AIDS Cases, Reported Through December 2002 | 34 |
| Table 6 | Cumulative Number of AIDS Cases, Reported Through December 2002, Selected Midwestern States and the U.S. | 35 |
| Table 7 | Prevalence for HIV by Age: July 2002 to June 2003 | 37 |
| Table 8 | Prevalence for AIDS by Age: July 2002 to June 2003 | 38 |
| Table 9 | Prevalence Rates of HIV/AIDS Infected Persons by Race/Ethnicity in 2003 | 40 |
| Table 10 | Prevalence Rates for HIV/AIDS by Mode of Transmission and by Race/Ethnicity: July 2002 to June 2003 | 43 |
| Table 11 | Prevalence Numbers for HIV and AIDS by Heterosexual Contact Categories | 44 |
| Table 12 | Estimated Number of People per Health Region in Indiana, 2002 | 45 |
| Table 13 | Prevalence Rates for HIV and AIDS by Health Region, July 2002 to June 2003 | 46 |
| Table 14 | Number of Infected Persons (HIV/AIDS) by State of Residence at the Time of this Report | 47 |
| Table 15 | Number of Infected Persons with HIV/AIDS by County of Residence at Time of Report | 48 |
| Table 16 | Number of New HIV/AIDS Cases, Reported in 2002 | 50 |
| Table 17 | Number of New HIV/AIDS Cases, Reported in 2002, Selected Midwestern States and the U.S. | 51 |
| Table 18 | Incidence Numbers for HIV, AIDS and HIV/AIDS by Sex, July 2002 to June 2003 | 53 |
| Table 19 | Incidence Numbers for HIV/AIDS by Race/Ethnicity, July 2002 to June 2003 | 55 |
| Table 20 | Incidence Numbers for HIV and AIDS, July 2002 to June 2003 | 56 |
| Table 21 | Incidence Numbers and Rates/100,000 for HIV/AIDS by Race/Ethnicity and Sex, July 2002 to June 2003 | 58 |
| Table 22 | Incidence Numbers, Percentages and Rates per 100,000 of HIV/AIDS | 59 |
| Table 23 | Incidence Rate for HIV/AIDS by Mode of Transmission and by Race/Ethnicity, July 2002 to June 2003 | 60 |
| Table 24 | Incidence Numbers for HIV/AIDS by Race/Ethnicity and Mode of Transmission, July 2002 to June 2003 | 60 |
| Table 25 | Incidence Numbers for HIV/AIDS by Indiana Health Region | 62 |

| | | |
|----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Table 26 | Incidence Numbers for HIV and AIDS by Health Region | 62 |
| Table 27 | Incidence Numbers for HIV, AIDS and HIV/AIDS by County | 63 |
| Table 28 | Cumulative Number of Children of HIV+ Mothers by the Mother's Infection Status and by the Diagnosis Status of the Child at the time of this Report, June 2003 | 67 |
| Table 29 | Cumulative Number of Children by their Infection Status and by the Availability of Drugs | 68 |
| Table 30 | Numbers of Persons Diagnosed with HIV or AIDS in Indiana and Currently Living outside the State | 73 |
| Table 31 | Number of Persons that were Diagnosed with HIV or AIDS outside Indiana and Migrated into Indiana | 74 |
| Table 32 | Cumulative Numbers of Migrants by Race and Ethnicity, 2003 | 76 |
| Table 33 | CTR Tests Performed by Gender in Indiana | 80 |
| Table 34 | CTR Tests Performed by Race and Ethnicity in Indiana | 80 |
| Table 35 | CTR Tests Performed by Age Group in Indiana | 81 |
| Table 36 | CTR Test Performed by Mode of Transmission in Indiana | 81 |
| Table 37 | CTR Tests Performed by Site Type in Indiana | 81 |
| Table 38 | CTR Tests Performed by Insurer of the Tested in Indiana | 82 |
| Table 39 | BRFSS Result: What was the main reason you had your last HIV test? | 84 |
| Table 40 | BRFSS Result: Where did you have your last HIV test? | 85 |
| Table 41 | STD Cases in Indiana, 1998 to 2003 | 88 |
| Table 42 | STD Cases in Indiana by Ethnicity, 2003 | 90 |
| Table 43 | STD Cases in Indiana by County, 2003 | 43 |
| Table 44 | Number of Pregnancies in Indiana by Age and Race, 2001 | 94 |
| Table 45 | Number of Pregnancies by County, 2001 | 95 |
| Table 46 | Number of Terminated Pregnancies in Indiana by County of Residence, 2001 | 97 |
| Table 47 | Title II Budget for Indiana, Fiscal Year 2003/2004 | 100 |
| Table 48 | Number of ADAP Recipients by Year and Race, 1999-2002 | 103 |
| Table 49 | Number of HIAP Recipients by Year and Race, 1999-2002 | 104 |
| Table 50 | Number of ADAP Recipients by Year and Ethnicity, 1999-2002 | 105 |
| Table 51 | Number of HIAP Recipients by Year and Ethnicity, 1999-2002 | 106 |
| Table 52 | Program Expenditures for ADAP by Year, 1999-2002 | 107 |

| | | |
|----------|--------------------------------------------------|-----|
| Table 53 | Program Expenditures for HIAP by Year, 1999-2002 | 108 |
| Table 54 | ADAP Pharmaceutical Activity, 1999 to 2002 | 108 |
| Table 55 | ADAP Clients on HAART, 2000-2002 | 108 |

Executive Summary

Demographics:

Indiana is a mostly rural state with several urban and metropolitan centers that had an estimated population of 6,159,068 people as of July 2002. The majority of the population (88.6%) is White and Non Hispanic, followed by African Americans (8.8%). The rest is comprised of people of Asian and Pacific Islander, and American Indian and Alaskan Native origin. The population is predominantly Non Hispanic (96.5%), with a small, but fast growing Hispanic minority. Based on the 2000 Census, 3.5% of the population selected Hispanic as their Ethnicity.

Prevalence:

By the end of June 2003, a total of 8087 people were infected with HIV/AIDS in the state of Indiana. The disease is male dominated, with the number of infected males five times higher than that of females. The rate of infection was at 219.2 for males and 44.2 for females per 100,000 people of the general population. Most of the infected persons are in their middle ages, ranging from 25 to 44 years of age. More than three out of ten people currently infected are African American, while about six out of ten people with HIV/AIDS were white. Based on the smaller numbers of African Americans in the general population, the infection rate of that racial group (531.5/100,000) is about three to six times the rate of the Hispanic (179.9/100,000) and White group (91.8/100,000). Three quarters of all infected males are White (52.8%) and African American (25.3%). Considering the much smaller number of African American males in the general population this demonstrates that African American males are much more affected compared to their White counterparts.

Each infected person is associated with a risk category of how they most likely were infected with the disease. The overwhelming majority of risk categories were Men Having Sex with Men (MSM). Its rate per 100,000 people of the population is between seven and ten times higher than the other risk categories for all infected people. It is the highest category of risk for all race and ethnicity groups, but it is especially pronounced for African Americans.

Geographically, the majority of infected people that were diagnosed in Indiana are also living here (95%). Within the state of Indiana, most infected people are concentrated in the urban areas of the state, with the majority living in Health region 6, corresponding to Central Indiana and the Indianapolis Metropolitan area, with 226.5 per 100,000 infected people. Other regions with large numbers of HIV/AIDS infected people include region 1 (125.0/100,000) which corresponds to the northwest part of the state adjacent to Chicago, and regions 5, 7 and 8 with about 90 per 100,000 of the general population.

Incidence:

The number of newly infected persons in Indiana was 593 for the period of July 2002 to June 2003. It had increased by almost 18% from the previous year and continues a trend of rising numbers of newly infected persons that started in the mid nineties. Consistent with the overall number of HIV/AIDS infected people, the majority of new infections is among males between the ages of 25 to 44 years of age. Males outrank females almost eight fold. Among the new infections with HIV/AIDS, African Americans have a rate that is twice as large as their Hispanic counterparts, and almost eight times that of Whites. Although African Americans make up only about 9% of the population, they account for almost 40% of all new infections. Consistent with the condition of the total infected population, males are predominant among new infections for all racial and ethnic groups. The rate of new infections with HIV/AIDS among African American males (68.0/100,000) is especially high, compared to their Hispanic (34.1) and White (10) counterparts. The highest rates are to be found in the MSM risk category, whose incidence rate of 8.2 per 100,000 is outranking all other risk categories and whose main contributors are African American (25.4/100,000) and Hispanic (16.2/100,000).

Geographically, over half of all newly infected persons live in Health Region 6 in Central Indiana, while the second largest group of almost 10% of all newly infected persons was living in region 1 in northwestern Indiana. Within the leading regions, Marion county and Lake county had the most new infections in the reported time period.

Mothers with HIV:

The number of reported cases of HIV positive mothers in Indiana was 50 in the reported time period. These mothers had a total of 421 children that were born, the majority after the mother tested positive for HIV. Almost two thirds of those mothers are African American, one quarter is White, and the rest were Hispanic women. Of all the children that were born to infected mothers, almost two third were definitely not infected, while 12% tested positive for HIV and 8% were diagnosed with AIDS. Please note that these numbers include all children, including those that were born before medication to prevent the spread of the HIV virus from mother to child was available.

Mortality:

The number of people that died of HIV/AIDS related complications in Indiana peaked around the year 1995 and started to drop sharply thanks to the widespread availability of anti-retroviral medications. However, the number has started to plateau in the last four years there were 137 deaths in 2002. The majority of infected people that died were White and African American males in the ages of 25 to 44. The majority of deaths are connected to the MSM risk group, with a mortality rate of 0.88 per 100 infected people that is more than three times that of the other rates. Geographically, the highest mortality rates occurred in regions 3 in Northeastern Indiana and zone 9 in East Central Indiana, with a rate of 3.1 and 3.4 death per 100 infected people respectively.

Mobility:

Of the total number of infected people, a relatively small number has migrated. At the end of June 2003, a total of 142 persons that were diagnosed with HIV/AIDS in Indiana and were still alive have moved out of the state, while at the same time a total of 505 people had moved to Indiana that were diagnosed with the disease in another state and that were alive at the time of this report. More infected White people have left the state than have moved here after they were diagnosed. In terms of numbers, 61.4% of infected persons that have left Indiana are White, compared to only 54.2% of those that moved to Indiana. In contrast, African Americans were represented in larger numbers among those that have left our state (38.0%) than those that moved here after they were diagnosed in another state (32.5%). A similar result is true for infected persons of Hispanic ethnicity. Among the population that moved to Indiana, their share (7.0%) of all infected people was smaller than that of the group moving out of the state (5.4%). Of those that moved to Indiana, more than a third have settled in central Indiana's Health Region 6. The rest is distributed more or less equally among the other regions of the state

In 2003, a total number of 19,010 HIV/AIDS tests were administered in Indiana, with 101 (or 0.4%) positive test results. The gender of the tested persons was almost equally split between males and females, while the rate of positive results for Whites, African American and Hispanic was about the same at 0.4% per 100 tested persons. The largest number of HIV positive results came from the 30 to 39 year old age range. Among the possible risk categories, the largest positive rate was among the MSM mode of transmission (2.0%).

Behavioral Risk Factor Surveillance System:

Assessing the indicators of risk for HIV/AIDS in Indiana, a survey was conducted that asked specific questions to a representative group of Indiana residents. The majority of Hoosiers, almost 96%, think that it is *Very Important* or *Somewhat Important* to know one's HIV status, but only about 40% have been tested before. Of those tests, the majority were done by a private doctor or HMO. Only half of the surveyed knew that an HIV positive pregnant woman can get treatment to help reduce the chances of passing the virus on to the child, while almost nine out of ten did have knowledge the availability of medical treatments that are intended to help a person who is infected with HIV to live longer. Finally, less than 10% of surveyed Hoosiers had a conversation with a health professional about preventing sexually transmitted diseases through condom use in the past 12 months.

STD:

Chlamydia is the most frequently reported sexually transmitted disease in Indiana, with 17,009 cases reported in 2003, followed by Gonorrhea with 6,656 cases and 379 cases of Syphilis. The numbers of new infections with Chlamydia and Gonorrhea have declined from their 2002 numbers, while Syphilis shows a slight increase compared to 2002. Females outnumber males for both Chlamydia and Gonorrhea

while Syphilis is more prevalent among males. Both African Americans and Whites make up the majority of all STD cases in the last year with an almost equal number of Chlamydia cases, while Gonorrhea and Syphilis are more widespread among African Americans. In 2002, Indiana had 85 cases of acute Hepatitis B and only one person with an acute infection with Hepatitis C. The total number of chronic Hepatitis C infections for the state was reported to be 6,313 people. And finally, 128 cases of TB were reported in Indiana in 2002, out of which 11 persons were also HIV positive.

Care Issues:

In the current fiscal year, that runs from April 1, 2003 to March 31, 2004, the funding for Title II of the Ryan White CARE Act added up to a total of \$10,555,376. The majority of that budget (74%) finances Health Insurance Assistance Programs (HIAP), while the rest is used for other medical and social services.

Of the 393 persons enrolled in the medical and social service programs called ADMS (AIDS Drug and Medical Services) in 2002, almost two thirds (63%) are White, and about a quarter (26%) are African American. The majority of recipients, 4 out of 10, reside in the Indianapolis Metropolitan area. The majority selected MSM as their main category of risk behavior. A total of 520 persons received assistance with the cost for medications in 2002 under the AIDS Drug Assistance Program (ADAP), while 1,364 clients were enrolled and received assistance through the Health Insurance Assistance Program (HIAP).

Unmet Needs are defined as service needs and gaps for infected individuals who know their HIV positive status and are not receiving regular primary medical care. In 2002, a total of 4726 persons had medical and service needs that were not met by the system. A final report of the findings about the Unmet Needs of infected persons in Indiana had not been issued at the time of this report.

Introduction

This statewide profile describes the epidemiology of HIV, AIDS and other sexually transmitted diseases (STD's) in Indiana, mainly through June 2003, the last date of surveillance information available. The report describes the distribution of the HIV/AIDS disease by geography, age, sex, race, ethnicity and associated risk categories.

The purpose of this profile is to assist in the development of a comprehensive HIV/AIDS prevention and care plan by providing the necessary data that can aid in designing and implementing prevention and care services, as well as pointing out the dimensions of unmet needs for infected persons throughout the state of Indiana.

In order to meet those criteria, the profile will address five key questions:

1. What are the sociodemographic characteristics of Indiana's population?
2. What is the scope of the HIV/AIDS epidemic in Indiana?
3. What are the indicators of risk for HIV/AIDS in Indiana?
4. What are the patterns of service utilization of HIV-infected people in Indiana?
5. What are the number and characteristics of people who know they are HIV-positive, but who are not receiving HIV primary medical care?

Profile Data Sources

This profile was compiled using a number of different data sources to present as complete as possible a picture of the epidemic. The majority of the information, however, will be derived from the HIV Surveillance Database, which was used heavily throughout the report.

Each data source has strength and limitations. A brief description of each source follows.

1. HIV/AIDS Surveillance

The data for this report was collected through 2002 and mid 2003 and includes reports of HIV infections, AIDS Indicator Diseases, HIV related laboratory results submitted by physicians, hospitals, counseling and testing sites, and laboratories.

2. Behavioral Surveys

Behavioral Risk Factor Surveillance System (BRFSS)

The BRFSS is a state-based random telephone survey of adults that collects data from Indiana residents on various modifiable health risks, preventive health measures, and demographic variables. The objective of the report is to encourage change in risk-related health behaviors, to discover target populations for programs and resources, and to present the condition of Indiana's health in a way that it can be compared to the health risk behaviors of the rest of the United States.

3. STD Surveillance Database

The STD Surveillance Database is part of the STD Program at the Indiana State Department of Health (ISDH) that provides technical and financial assistance to local STD Programs for surveillance, case detection through screenings, ensuring treatment of known cases, case follow-up and education. Efforts are coordinated among the groups screening for Syphilis, Gonorrhea, and Chlamydia. The data in this database comes from STD clinics, physicians, and laboratories for STD infections (Syphilis, Gonorrhea, Syphilis) diagnosed through 2003.

4. Counseling, Testing and Referral Data

This database is compiled from the statewide outreach program that coordinates the efforts of local HIV counseling and testing sites. The effort is intended to promote the early detection of HIV infection and the management of HIV-related illnesses, educate people regarding alternative behaviors to those which facilitate HIV infection, and to facilitate access to health care. The data that is collected at Counseling, Testing and Referral sites throughout Indiana. Testing is voluntary and either anonymous or confidential. The tests include either blood or oral testing medium.

5. Vital Statistics

The Vital Statistics source provides excerpts from the Natality report and pregnancy outcomes for 2001.

6. Population Data

The main source for all population data in the U.S. is the Census Bureau that conducts a general population survey every ten years. The last decennial census was conducted in 2000. The results of that census, as well as forecasts and estimates for 2002 were issued by the Census Bureau and are used in this Epidemiologic Profile.

7. HIV Care Data

The HIV Care Data is collected through the HIV Care Coordination Program that offers a specialized form of case management and is the foundation upon which all other HIV/AIDS services are built. The service is available statewide through seventeen (17) regional sites which provide HIV positive residents with an individualized plan of care to address their medical, psychosocial, financial, and other supportive services needs.

Strengths and Limitations of Profile

LIMITATIONS AND CAVEATS

Data collected through the Indiana HIV/AIDS surveillance system is collected through: 1) required reports of persons with HIV infection and AIDS diagnosed or treated by physicians and hospitals, and 2) by laboratories that test for HIV, indicators for HIV, an antigen of the virus, or antibodies to that virus. Indiana reporting is designed such that the surveillance system receives multiple reports on each diagnosed person as the disease progresses and as various health care providers serve the person. AIDS case reports are the only HIV-related data consistently available on a population-wide basis in all states by sex, race/ethnicity, age, and mode of HIV exposure (risk factor). Job Corps entrants and civilian applicants for military service are also tested for HIV and reported to the CDC as an aggregate.

HIV and AIDS case report data represents those persons who: 1) have confidentially tested positive for HIV, 2) received medical care, and 3) have been reported by the health care provider. Deaths are voluntarily reported and gleaned from death certificates with HIV and AIDS identified as a cause of death.

Data from the federally funded and state administered HIV Counseling and Testing Services (CTS) program's counseling and testing sites may assist in assessing the characteristics of newer HIV infections except that CTS clinic data have several limitations. CTS clients are self selected and do not necessarily represent more recent infections in the state. The data represent tests performed, not persons tested, and the tests include unlinked results and repeated tests. Those tests that are performed confidentially and test positive for HIV are reported to the surveillance system by name and thus are linked and unduplicated in the surveillance reporting system. Comparing CTS data based on tests done and surveillance data based on reported cases has some variation in all groups.

Data are also included from the Sexually Transmitted Disease program, Tuberculosis Control program, Behavior Risk Factor Surveillance program, and the Vital Statistics Unit at the Indiana State Department of Health. Data were also included from the Center's for Disease Control and Prevention's HIV Surveillance report.

It is advisable not to over-interpret small changes or differences from year to year or between different groupings within the same year. Given the low numbers of cases in some categories, they may be misleading. Also misleading can be slight differences in exact numbers, depending on which database they were derived. In those cases it is more the general tendency that matters and not the exact absolute number. All data sources are not equivalent, and not all databases will add up to the exact same numbers.

Organization of the Profile

This profile answers five core questions in two main sections.

Chapter 1: Core Epidemiologic Questions

This section provides background on the characteristics of the general population of Indiana, distribution of HIV, and a profile of those at risk for HIV infection. Three questions form the core of this section:

1. **What are the sociodemographic characteristics of the general population in Indiana?** Informs readers about the overall demographic and socioeconomic characteristics of Indiana.
2. **What is the scope of the HIV/AIDS Epidemic in Indiana?** Assesses the impact of HIV/AIDS among a number of groups within Indiana. This allows planners and care givers to target those populations most at risk.
3. **What are the indicators of risk for HIV/AIDS in Indiana?** Provides an in-depth look at populations most at risk. Indirect and direct measures assist in indicating high-risk behaviors.

Chapter 2: Ryan White HIV/AIDS CARE Act Special Questions and Considerations

This section concentrates on questions pertaining to Health Resources and Services Administration HIV/AIDS care planning. Included is a description of access, utilization, and standards of care among HIV-positive people in Indiana. It is organized around two central questions:

1. **What are the patterns of service utilization of HIV-infected people in Indiana?** Examines service utilization by several populations living with HIV/AIDS.
2. **What are the number and characteristics of people who know they are HIV-positive, but who are not receiving HIV primary medical care?** Estimates unmet need of persons who are HIV-positive, but not receiving care throughout Indiana.